FIFRA 2(ee) Recommendation

EPA Reg. No. 279-9720

FOR DISTRIBUTION AND USE ONLY IN:
Colorado, Kansas, Illinois, Indiana, Missouri, Nebraska, North Dakota, Oklahoma, and South Dakota

THIS RECOMMENDATION, WHICH CONTAINS ADDITIONAL DIRECTIONS FOR USE, IS MADE AS PERMITTED UNDER FIFRA SECTION 2(ee) AND HAS NOT BEEN SUBMITTED TO OR APPROVED BY THE EPA.

This recommendation for STEWARD eVo Insect Control is valid until 12/31/2031, or until withdrawn, canceled or suspended.

IT IS A VIOLATION OF FEDERAL LAW TO USE THIS PRODUCT IN A MANNER INCONSISTENT WITH ITS LABELING. ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA REGISTERED LABEL MUST BE FOLLOWED.

THESE USE DIRECTIONS MUST BE IN THE POSSESSION OF THE USER AT THE TIME OF PESTICIDE APPLICATION.

FOR USE OF STEWARD eVo INSECT CONTROL FOR CORN LEAFHOPPER ON CORN

CROP	PEST	RATE OF APPLICATION
Corn (field) Corn (grown for seed) Corn (pop)	• Corn Leafhopper (*Dalbulus maidis)*	3.0 – 5.6 fl. oz./A

APPLICATION INSTRUCTIONS AND RESTRICTIONS:

- The minimum interval between treatments is 5 days.
- Make no more than 2 applications per acre per crop.
- Do not apply more than 11.25 fl oz/A of STEWARD eVo Insect Control or 0.22 lb ai/A of indoxacarb-containing products per calendar year.
- Apply in a minimum of 3 GPA (aerial) and 10 GPA (ground) for optimal coverage and efficacy.
- Application by chemigation systems is allowed.
- The REI is 12 hours.
- Interval between last application and harvest is: 1 day for forage, fodder, and silage (field and corn
- grown for seed only) and 14 days for grain and stover.

STEWARD eVo Insect Control may be mixed with adjuvants, fungicides, or herbicides to provide disease or weed control and insect management in one application.

R-5061 090426

FMC Corporation
2929 Walnut St.
Philadelphia, PA 19104
215-299-6000

STEWARD and FMC – Trademarks of FMC or an affiliate
©2026 FMC Corporation. All rights reserved.